

CHAPTER 2

THE PAIRING THEORY OF SUPERCONDUCTIVITY

In analogy with a free electron gas, normal (N) metals exhibit a single-particle excitation spectrum which, in the limit of a large system, is a continuum starting at zero energy. The degeneracy associated with this spectrum leads to the linear temperature dependence of the electronic specific heat near 0°K , and to the large electrical and thermal conductivities of these materials. In the superconducting (S) phase, the single-particle excitation is radically different from that of normal metals. In superconductors a minimum energy 2Δ , called the energy gap, is required to make a single-particle excitation from the ground state.

2-1 PHYSICAL NATURE OF THE SUPERCONDUCTING STATE

This qualitative difference in the excitation spectra is paralleled by a qualitative difference in the wave functions of the N - and S -phases of metals. In the N -phase, the probability that two single-particle states i and j are simultaneously occupied

is a smoothly varying function of the quantum numbers i and j . For example, in a pure single crystal, the expectation value

$$P_{kk'}^N = \langle N | n_{k\uparrow} n_{k'\downarrow} | N \rangle \quad (2-1)$$

is a smoothly varying function of \mathbf{k} and \mathbf{k}' (so long as one does not cross the Fermi surface in varying \mathbf{k} or \mathbf{k}'). Here $|N\rangle$ represents a typical state in the normal phase and $n_{k\uparrow}$ is the operator which measures the number of electrons in $\mathbf{k}\uparrow$, etc. (see the Appendix). In the superconducting phase,⁸ the corresponding probability

$$P_{kk'}^S = \langle S | n_{k\uparrow} n_{k'\downarrow} | S \rangle \quad (2-2)$$

is also a smoothly varying function of \mathbf{k} and \mathbf{k}' except when \mathbf{k} and \mathbf{k}' are related by the "pairing" condition. This condition states that for a given state \mathbf{k} , there exists a single mate $\bar{\mathbf{k}}$ such that the probability $P_{k\bar{k}}^S$ is larger than $P_{kk'}^S$ by a finite amount, for all states \mathbf{k}' in the vicinity of $\bar{\mathbf{k}}$. This singular behavior of the two-particle correlation function, which has been stressed by Yang,³⁹ is no doubt the sort of picture F. London had in mind when he suggested that superconductivity is due to a condensation of the electrons in momentum space.¹ When proper account is taken of residual interactions conventionally neglected in the description of the normal state, these "pairing correlations" leading to superconductivity emerge in a natural manner. Above the superconducting transition temperature, the pairing correlations are broken up by thermal fluctuations and play no important role in the normal phase.

It is essential to realize at the outset that the lowering in energy of the S -phase due to interactions between mates (say $\mathbf{k}\uparrow$ and $\bar{\mathbf{k}}\downarrow$) of a given pair depends critically on the choice of mates ($\mathbf{k}'\uparrow$ and $\bar{\mathbf{k}}'\downarrow$) for other pairs. In fact, the energy gap and most of the observed properties of the superconducting phase would be *absent* were it not for strong correlations *between the pairs*. The reason for the simple BCS model working so well is that in real metals these pair-pair correlations are almost entirely due to Pauli principle restrictions rather than correlations due to true dynamical interactions between the pairs. This fact allows one to treat the system in lowest order as if dynamical interactions

existed *only between mates* of a pair. The pair-pair correlations would then be accounted for by working out this reduced problem consistently with Fermi-Dirac statistics so as to include the crucial Pauli principle correlations between the pairs. We shall call this scheme the pairing (or BCS) approximation.

For a translationally invariant system, we shall see that the pairing $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$ of Bloch states leads to the lowest energy of the system. Supercurrent-carrying states are generated by translating this state of the system by an amount $\mathbf{q}/2$ in \mathbf{k} -space. The pairing would then be $(\mathbf{k} + \mathbf{q}/2 \uparrow, -\mathbf{k} + \mathbf{q}/2 \downarrow)$ and the electrons would have a net drift velocity $\mathbf{v}_d = \hbar\mathbf{q}/2m$. More generally, corresponding to each physical system and each state of that system there is a choice of pairing of single-particle states which minimizes the energy (or free energy, at finite temperature). For example, in a superconductor with nonmagnetic impurities present one should pair one-electron states φ_n which include the impurity scattering potential, as Anderson⁴⁰ first pointed out. He showed that one should pair a state φ_n and its time-reversed mate φ_n^* to form the ground state of the system in this case. For a uniform hollow cylinder in the absence of a magnetic field, one would pair the state (n, m, k) with its time-reversed partner $(n, -m, -k)$, where n and m are the radial and azimuthal quantum numbers, respectively, and k is the wave number for motion along the axis of the cylinder. In the presence of a magnetic field, the best pairing depends on the thickness of the cylinder and the strength of the field. For a thickness $d \gg \lambda$ (the penetration depth) one would pair $(n, m + \nu, k)$ with $(n, -m + \nu, k)$ or $(n, m + \nu, k)$ with $(n, -m + \nu + 1, k)$ depending on whether the flux trapped in the hole is an even or odd multiple of the flux quantum $\hbar c/2e$, that is, $\nu\hbar c/e$ or $(\nu + \frac{1}{2})\hbar c/e$. As we shall see, these are the only allowed values of the flux trapped within a thick-walled superconducting cylinder. We shall study in detail these various possible pairings in later chapters when we apply the basic theory to physical problems.

While the above "pairing" approximation gives a good account of the single-particle excitation spectrum, there exist

collective modes, such as the plasmons, arising from residual interactions neglected within this approximation. In addition there may be small momentum exciton-like collective modes which lie within the gap. For larger momentum, the exciton states rise above the gap edge and pass into the continuum, thereby becoming heavily damped. The nature of the collective states and their effect on system properties is discussed in Chapter 8.

Also neglected within the simplest pairing approximation are damping effects. In the strong-coupling superconductors, such as lead and mercury, it is essential to include these effects on the same footing as the pairing correlations to obtain a reasonable description of these "bad actors."

The above discussion suggests that the excited states of a superconductor can be represented by a two-fluid model, one for the condensed electrons and one for the excitations. As we mentioned in the introduction, phenomenological two-fluid models (most notably the Gorter-Casimir model² and the Ginsburg-Landau model³⁶) have played an extremely important role in laying the ground work for our present understanding of superconductivity. While there are important differences between the predictions of the pairing theory and the earlier two-fluid models, the theories share the basic idea that the superfluid electrons (i.e., the strongly correlated pairs in our case) can be described by a local density $\rho_s(\mathbf{r})$ and a local flow velocity $\mathbf{v}_s(\mathbf{r})$. The excited electrons then form an interpenetrating normal fluid which in local thermal equilibrium can be described by the local quantities $\rho_n(\mathbf{r})$ and $\mathbf{v}_n(\mathbf{r})$. As we shall see, the superfluid can only carry out potential flow, that is $\text{curl } \mathbf{v}_s(\mathbf{r}) = 0$, a condition emphasized by F. London.¹ (No such restriction holds for the normal fluid.) Many of the observed properties of superconductors can be understood in terms of a two-fluid model having a temperature-dependent energy gap for creating normal fluid (excitations) from the superfluid component.^{16c} As we shall see, one can often interpret the results of the microscopic theory in terms of such a model.⁹

2-2 THE ONE-PAIR PROBLEM

To understand the origin and consequence of pairing correlations, it is helpful to consider the problem, first studied by Cooper,⁴¹ of a pair of electrons interacting above a noninteracting Fermi sea of electrons via a velocity-dependent nonretarded two-body potential V . Thus, all but two of the electrons are assumed to be noninteracting. The background electrons enter the total problem only through the Pauli principle by blocking states below the Fermi surface from participating in the remaining two-particle problem. If we measure the kinetic energy ϵ_k relative to its value at the Fermi surface, only states with $\epsilon_k > 0$ are available to the interacting pair of electrons. Since the system is assumed to be translationally invariant and one neglects spin-dependent forces, the center-of-mass momentum $\hbar\mathbf{q}$ of the pair and the total spin S are constants of motion. The orbital wave function of the pair can then be written as

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \varphi_{\mathbf{q}}(\boldsymbol{\rho})e^{i\mathbf{q}\cdot\mathbf{R}} \quad (2-3)$$

where the relative and center-of-mass coordinates are defined by $\boldsymbol{\rho} = \mathbf{r}_1 - \mathbf{r}_2$ and $\mathbf{R} = (\mathbf{r}_1 + \mathbf{r}_2)/2$, respectively. The relative coordinate wave function is symmetric for the singlet spin state ($S = 0$) and antisymmetric for the triplet states ($S = 1$). In the limit $\mathbf{q} \rightarrow 0$ the relative coordinate problem is spherically symmetric so that $\varphi(\boldsymbol{\rho})$ is an eigenfunction of angular momentum and can be labeled by the angular momentum quantum numbers l and m . For $\mathbf{q} \neq 0$, the component of angular momentum along \mathbf{q} and parity remain good quantum numbers but l is no longer sharp.

For simplicity, we first consider the zero momentum states $\mathbf{q} = 0$, so that ψ can be expanded as

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \varphi(\boldsymbol{\rho}) = \sum_{\mathbf{k}} a_{\mathbf{k}} e^{i\mathbf{k}\cdot\boldsymbol{\rho}} = \sum_{\mathbf{k}} a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}_1} e^{-i\mathbf{k}\cdot\mathbf{r}_2} \quad (2-4)$$

In (2-4), the sum is restricted to the available states ($\epsilon_k > 0$). Since the factors $e^{i\mathbf{k}\cdot\mathbf{r}_1}$ and $e^{-i\mathbf{k}\cdot\mathbf{r}_2}$ can be thought of as single-particle states of momentum \mathbf{k} and $-\mathbf{k}$, we see that the pair wave function is a superposition of configurations in each of which a definite pair state $(\mathbf{k}, -\mathbf{k})$ is occupied.

To find the zero-spin eigenstates of the pair, we write Schrödinger's equation as

$$(W - H_0)\psi = V\psi \quad (2-5a)$$

and from (2-4) one has

$$(W - 2\epsilon_k)a_k = \sum_{k'} V_{kk'}a_{k'} \quad (2-5b)$$

where the matrix element $V_{kk'}$ is defined by

$$V_{kk'} = \langle \mathbf{k}, -\mathbf{k} | V | \mathbf{k}', -\mathbf{k}' \rangle \quad (2-6)$$

In Figure 2-1 a typical scattering process caused by V is illustrated.

While the Schrödinger equation (2-5b) cannot be analytically solved in general, the solution is immediate if $V_{kk'}$ is taken to be a factorizable potential $V_{kk'} = \lambda w_{k'}^* w_k$. More generally, if the system is isotropic $V_{kk'}$ can be expanded into its partial wave components

$$V_{kk'} = \sum_{l=0}^{\infty} \sum_{m=-l}^l V_l(|\mathbf{k}|, |\mathbf{k}'|) Y_l^m(\Omega_k) Y_l^{-m}(\Omega_{k'}) \quad (2-7)$$




FIGURE 2-1 A typical transition occurring in Cooper's problem in which one pair of electrons interacts above a quiescent Fermi sea. The center-of-mass momentum of the pair is chosen to be zero in this drawing.

and the (l, m) eigenstates of the pair can be determined if V_l is taken to be factorizable,

$$V_l(|\mathbf{k}|, |\mathbf{k}'|) = \lambda_l w_k^l w_{k'}^{l*} \quad (2-8)$$

In this case we have from (2-5b)

$$(W_{lm} - 2\epsilon_k) a_k = \lambda_l w_k^l \sum_{k'} w_{k'}^{l*} a_{k'} \quad (2-9a)$$

where

$$a_k = a_k Y_l^m(\Omega_k) \quad (2-9b)$$

Equation (2-9a) can be written as

$$a_k = \frac{\lambda_l w_k^l C}{W_{lm} - 2\epsilon_k} \quad (2-10a)$$

where the constant C is defined as

$$C = \sum_{k'} w_{k'}^{l*} a_{k'} \quad (2-10b)$$

By substituting (2-10a) into the definition (2-10b) one obtains the equation

$$1 = \lambda_l \sum_k |w_k^l|^2 \frac{1}{W_{lm} - 2\epsilon_k} \equiv \lambda_l \Phi(W_{lm}) \quad (2-11)$$

determining the energy eigenvalues W_{lm} . If we work in a large but finite box the single-particle energies ϵ_k form a discrete set so that when W passes from below to above $2\epsilon_k$, $\Phi(W)$ jumps from $-\infty$ to ∞ . As W moves toward the next higher value of $2\epsilon_k$, $\Phi(W)$ again approaches $-\infty$ and jumps to $+\infty$ as W passes through this higher value. The function $\Phi(W)$ is shown schematically in Figure 2-2. As W passes through the origin to negative values (i.e., the region of bound states) $\Phi(W)$ increases from $-\infty$ to zero as shown. The eigenvalues W_{lm} are given by the intersections of $\Phi(W)$ with the constant function $1/\lambda_l$, as shown for both positive (repulsive) and negative (attractive) λ_l . While the eigenvalues in the continuum are trapped between the unperturbed energies $2\epsilon_k$ and approach the unperturbed energies as the size of the box goes to infinity, a state is bound

split off from the continuum for an attractive l -wave potential, as shown in the figure. For the simple case

$$w_k^l = \begin{cases} 1 & 0 < \epsilon_k < \omega_c \\ 0 & \text{otherwise} \end{cases} \quad (2-12a)$$

and $\lambda_l < 0$, the binding energy $|W_{lm}|$ of the pair in the split-off state is given by

$$\frac{1}{|\lambda_l|} = \frac{N(0)}{2} \log \left[\frac{|W_{lm}| + 2\omega_c}{|W_{lm}|} \right] \quad (2-12b)$$

or

$$|W_{lm}| = \frac{2\omega_c}{\exp \left[\frac{2}{N(0)|\lambda_l|} \right] - 1} \quad (2-12c)$$

We have assumed the density of states $N(\epsilon_k)$ is slowly varying in the interval $0 < \epsilon_k < \omega_c$ and have approximated it by $N(0)$, the density of single-electron states of one-spin orientation, evaluated at the Fermi surface. From (2-12c) one has for weak coupling [$N(0)|\lambda_l| \ll 1$]

$$|W_{lm}| \cong 2\omega_c \exp \left[-\frac{2}{N(0)|\lambda_l|} \right] \quad (2-13a)$$

while for strong coupling [$N(0)|\lambda_l| \gg 1$] one obtains

$$|W_{lm}| \simeq N(0)|\lambda_l|\omega_c \quad (2-13b)$$




FIGURE 2-2 A plot of the function $\Phi(W)$ [see (2-11)] which determines the eigenenergies in Cooper's one-pair problem. For a repulsive interaction ($\lambda_l > 0$), all states are trapped in the continuum, while for an attractive interaction, a bound state is split off.

From (5-13a) we see that the binding energy is an extremely sensitive function of the coupling strength for weak coupling; however, a bound state exists for arbitrarily weak coupling so long as the potential is attractive near the Fermi surface. This important result was discovered by Cooper,⁴¹ who suggested that the instability of the normal phase, because of pairs of electrons entering this type of bound state, was associated with the occurrence of the superconducting phase.

Earlier work of Schafroth, Blatt, and Butler (SBB)⁴² is closely related to Cooper's discussion. Schafroth⁴³ had suggested that the superconducting state might correspond to a Bose-Einstein condensation of pairs of electrons into localized bound states. An attempt to develop a theory along these lines was made by SBB using what they call the quasi-chemical equilibrium approach for evaluating the partition function of the system. Owing to mathematical difficulties, they were not able to carry out calculations based on their general formulation for any model which exhibited superconducting properties. For a qualitative picture they suggested a model with localized pairs such that the size of the pair bound state is small compared to the average spacing between pairs. The bound pairs would presumably be capable of translational motion relative to the other pairs and one would obtain a continuum of Bose-Einstein excitations above the ground state without an energy gap, in contrast with the pairing theory. If the pairs were indeed well separated they could be treated as independent and Cooper's discussion would be appropriate. It should be pointed out, however, that, subsequent to the work of Bardeen, Cooper, and the author, Blatt and Matsubara extended the Bose condensation approach to give the results of the pairing theory.

Actual superconductors differ in a fundamental manner from a bound pair model in which the pairs are either well separated in space and/or weakly interacting. As we shall see below, there are on the average about one million bound pairs which have their centers of mass falling within the extent of a given pair function. Thus, rather than weakly overlapping pairs, one has just the reverse

limit, very *strongly* overlapping pairs. As we mentioned above, it is surprising that one can meaningfully treat such a system in zero order by including only dynamical interactions between mates of a pair and neglect all but the Pauli principle restrictions when treating interactions between the pairs. It is intended that the discussion below and in the following chapters will help clarify this point.

Returning to the one-pair problem of Cooper, it is interesting to see how the energy of the bound state varies with the center-of-mass momentum $\hbar\mathbf{q}$. If we assume only the *s*-wave ($l = 0$) part of V is important (as appears to be the case except for the crystalline anisotropy effects), one finds that the binding energy $W(\mathbf{q})$ satisfies

$$1 = |\lambda_0| \sum_{\mathbf{k}} \frac{1}{|W_{\mathbf{q}}| - \epsilon_{\mathbf{k}+\mathbf{q}/2} - \epsilon_{\mathbf{k}-\mathbf{q}/2}} \quad (2-14)$$

where $|\mathbf{k} + \mathbf{q}/2|$ and $|\mathbf{k} - \mathbf{q}/2|$ are required to be greater than k_F , and the sum is to be cut off at $\epsilon_{\mathbf{k}} = \omega_c$. For small q one finds

$$|W_{\mathbf{q}}| = |W_0| - \frac{v_F \hbar q}{2} \quad (2-15)$$

where

$$|W_0| = \frac{2\omega_c}{\left\{ \exp \left[\frac{2}{N(0)|\lambda_l|} \right] - 1 \right\}}$$

as above. Thus, the pair energy increases linearly with the center-of-mass momentum in the limit $q \rightarrow 0$, rather than as q^2 , as one might expect. As Cooper pointed out, the drift of the pair with respect to the noninteracting Fermi sea strongly reduces the binding energy of the pair due to the reduced density of low-energy states available to the pair. This effect dominates the q^2 increase of kinetic energy for small q .

If $|W_0|$ is imagined to be of order kT_c , (2-15) shows that the pair would have lost most of its binding energy when

$$q \sim \frac{k_B T_c}{\hbar v_F} \sim \frac{k_B T_c}{E_F} k_F \sim 10^{-4} k_F \sim 10^4 \text{ cm}^{-1} \quad (2-16)$$

This number is roughly numerically equal to the reciprocal of

Pippard's coherence length $\xi_0 \sim 10^{-4}$ cm,³³ about which we shall have more to say. If one calculates the pair function $\varphi(\rho)$ from (2-4) and (2-10a) one finds the size of the bound state is of order ξ_0 . Thus, one would be required to have an extremely small density of bound pairs if an isolated pair model were to be appropriate. In fact, the density would be so small that the predicted N - S energy difference at zero temperature would be many orders of magnitude too small to agree with experiment.

Thus far we have considered only the singlet spin state of the bound pair. If there is a strong attractive odd l potential, the triplet state will have the largest binding energy and one might expect the pairing in the superconducting state to be in a triplet state. There is no experimental evidence to support other than singlet pairing at present.

In closing this section we emphasize that the single-pair model exhibits a continuous spectrum above the ground state, without an energy gap.

2-3 LANDAU'S THEORY OF A FERMI LIQUID

Looking back at Cooper's argument, one might raise several objections to the conclusion that the bound state in the two-particle problem has anything to do with the occurrence of superconductivity. For example, one knows that Coulomb and phonon interactions between electrons in the normal state lead to a correlation energy of order one electron-volt per electron,⁴⁴ compared with the negligibly small binding energy $W \simeq 10^{-4}$ ev of a bound pair. Is it not possible that the strong correlations between all the electrons in the normal state will necessarily lead to fluctuations which break up the weakly bound state of a given pair of electrons? In addition, even if such a bound state could exist in a metal, would not the strong overlap of the pairs required to fit the observed condensation energy lead to interactions which would destroy the concept of bound pairs?

In answering the first objection, it is important to recognize that Landau's theory of a Fermi liquid⁴⁵ gives a good account of the low-lying single-particle excitations of the normal state. In

this theory, the excited states of a normal metal are placed in one-to-one correspondence with those of a free-electron gas. In Landau's theory, one asserts that the essential effect of the interactions between the electrons in the normal state of a metal is to shift the effective mass of an electron (now called a "quasi-particle") by an amount which is observed to be of order 10 to 50 per cent. An important feature of the theory is that a quasi-particle, as opposed to a "bare" electron, is a stable excitation in the immediate vicinity of the Fermi surface (at sufficiently low temperature). There is, however, a coupling between quasi-particles which arises from interactions neglected within this Fermi liquid approximation. This residual coupling leads to superconductivity.

The basis for Landau's theory has been extensively investigated and one knows that the theory is correct in all orders of perturbation theory⁴⁶ starting from the noninteracting system. The theory no doubt has a wider range of validity than that of the perturbation series itself, although its exact limitations are not known at present. Empirically, Landau's theory works very well in the normal state.

The remarkably small energy difference between the normal and superconducting states of a metal (10^{-8} ev per electron) strongly suggests that there is only a subtle shift of the electron-electron correlations between the two states. Since Landau's theory gives a good account of the normal state, it is reasonable to use the complete set of wave functions given by this theory as a basis for constructing the wave functions of the superconducting state. This procedure is particularly appealing because the superconducting wave functions primarily involve normal state configurations in which quasi-particle excitations are present only near the Fermi surface. However, these are just the configurations which are best described by the Landau theory. Therefore, Cooper's result is to be understood in the sense that his two-particle problem is actually a two-quasi-particle problem.

A difficulty with the above approach is that one knows little about the interaction between quasi-particles in the normal phase

from an experimental point of view. While the quasi-particle's effective mass involves the forward scattering amplitude of two quasi-particles, the effective mass also involves band-structure effects which are difficult to estimate accurately. More important is the fact that one needs the interaction for finite momentum and energy transfer so that one is forced to estimate the interaction theoretically. While this problem is not completely settled at present, it appears that one understands the essential features of the interaction, the remaining complications arising primarily from detailed crystalline effects (see Chapter 7).

In regard to the second objection mentioned above, it is true that the simple picture of bound pairs of electrons forming the ground state of the superconductor is impaired by their great overlap. Nevertheless there remains the strongly correlated occupancy of a given quasi-particle state (say $\mathbf{k} \uparrow$) with its mate (say $-\mathbf{k} \downarrow$) as in Cooper's problem. Interactions between quasi-particles which tend to break up this correlated occupancy are presumably already included in Landau's description of the normal state. Thus a simplified model in which one includes pairing correlations between otherwise noninteracting quasi-particles is not at the outset an unreasonable starting point. It is this point of view Bardeen, Cooper, and the author took in constructing the microscopic theory of superconductivity.

2-4 THE PAIRING APPROXIMATION

We saw above that for a translationally invariant normal system carrying no current, the $\mathbf{q} = 0$ pair state is the most unstable, in the sense that it is the pair with the largest binding energy W . In Chapter 7 a time-dependent treatment of the normal state instability is given and one finds the greatest growth rate is for $\mathbf{q} = 0$ pairs, in agreement with Cooper's result. Thus it is natural to solve the reduced problem in which interactions are considered only between electrons of opposite momentum. One hopes that the resolution of the strongest instability will modify the system so as to remove the $\mathbf{q} \neq 0$ pair instabilities as well. This is exactly what happens. We restrict our attention to pairing electrons of opposite spin orientation.

We shall use the formalism of second quantization to describe the interacting electron system; this scheme is reviewed in the Appendix. The creation and destruction operators for electrons of wavevector \mathbf{k} and z -component of spin s (\uparrow or \downarrow) are denoted by c_{ks}^+ and c_{ks} , respectively. They satisfy the usual Fermi anticommutation rules. The Hamiltonian for the reduced problem of the $\mathbf{q} = 0$ pairs is

$$H_{\text{red}} = \sum_{ks} \epsilon_k n_{ks} + \sum_{k'k} V_{k'k} b_{k'}^+ b_k \quad (2-17)$$

where the pairing matrix element $V_{k'k}$ is given by

$$V_{k'k} = \langle k', -k' | V | k, -k \rangle \quad (2-18)$$

and the operator b_k^+ creates a pair of electrons in the single-particle states $k \uparrow$ and $-k \downarrow$, that is,

$$\begin{aligned} b_k^+ &= c_{k\uparrow}^+ c_{-k\downarrow}^+ \\ b_k &= c_{-k\downarrow} c_{k\uparrow} \end{aligned} \quad (2-19)$$

This type of Hamiltonian forms the basis of the theory of superconductivity proposed by Bardeen, Cooper, and the author.⁸ Further argument for concentrating on these particular interactions in describing the ground state and the low-lying excited states are given in the original paper by Bardeen, Cooper, and the author and in a review article by Bardeen and the author.⁹ There it is argued, on the basis of phase-space considerations as well as effects due to the antisymmetry of the wave functions, that the $\mathbf{q} = 0$ pair state should be macroscopically occupied in cases where the superfluid momentum density is zero (although \mathbf{v}_s need not be zero in the presence of magnetic fields). We expect the matrix element $V_{k'k}$ to be predominantly negative near the Fermi surface for superconductivity to occur. As we shall see in Chapter 7, this attraction is due to the ions overscreening the Coulomb repulsion, thereby reversing the sign of the effective interaction. While (2-17) is written in terms of bare single-particle operators c_{ks} , the reduced Hamiltonian can be formally viewed as a model Hamiltonian describing residual interactions ($V_{k'k}$) between the quasi-particles in the normal phase as discussed above. Since the pairing correlations constitute a fractional change of only $\sim 10^{-8}$ in the total correlation energy of a metal, it is clear that this more

liberal interpretation of H_{red} must be adopted. When the quasi-particle picture of the normal phase excitations is insufficient, as in lead and mercury, other techniques must be used in treating the superconducting phase (see Chapters 6 and 7). Note that the form of H_{red} depends upon the choice of pairing, as discussed in Section 2-1.

Since the pairing interaction maintains the pairing condition, it follows that the eigenstates of H_{red} can be labeled by those states \mathbf{k}, s which are occupied, their mates $-\mathbf{k}, -s$ being unoccupied. This labeling of states leads to a one-to-one correspondence of the eigenstates of H_{red} with the eigenstates of a noninteracting Fermi gas (or the normal state). As we shall see, if excitations happen to be in states $\mathbf{k} \uparrow$ and $-\mathbf{k} \downarrow$, special care must be taken so that these configurations are properly orthogonal to the ground-state wave function.

It is clear that if V is attractive, the ground state of H_{red} has no pair state $(\mathbf{k} \uparrow, -\mathbf{k} \downarrow)$ occupied by a single electron. In this case the operator $n_{\mathbf{k} \uparrow} + n_{-\mathbf{k} \downarrow}$ can be replaced by $2b_{\mathbf{k}}^+ b_{\mathbf{k}}$ that is, twice the pair occupation number. The reduced Hamiltonian is then

$$H_{\text{red}}^0 = \sum_{\mathbf{k}} 2\epsilon_{\mathbf{k}} b_{\mathbf{k}}^+ b_{\mathbf{k}} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}'\mathbf{k}} b_{\mathbf{k}'}^+ b_{\mathbf{k}} \quad (2-20)$$

It might be argued that eigenstates of H_{red}^0 follow immediately by forming new operators B_n which are linear combinations of the $b_{\mathbf{k}}$'s such that H_{red} is of the form $\sum_n \tilde{\epsilon}_n B_n^+ B_n$. This argument is incorrect. If the operators $b_{\mathbf{k}}$ and $b_{\mathbf{k}}^+$ described true Bose particles (rather than pairs of fermions) the B_n 's and B_n^+ 's would also describe Bose particles and the ground state would be formed by placing all the bosons in the lowest state. Rather, one finds by direct calculation the commutation relations

$$[b_{\mathbf{k}}, b_{\mathbf{k}'}^+] = 0 \quad \text{for } \mathbf{k} \neq \mathbf{k}' \quad (2-21a)$$

$$[b_{\mathbf{k}}, b_{\mathbf{k}'}^+] = 1 - (n_{\mathbf{k} \uparrow} + n_{-\mathbf{k} \downarrow}) \quad \text{for } \mathbf{k} = \mathbf{k}' \quad (2-21b)$$

and

$$[b_{\mathbf{k}}, b_{\mathbf{k}'}] = 0 = [b_{\mathbf{k}}^+, b_{\mathbf{k}'}^+] \quad (2-21c)$$

These are not of the form required by Bose-Einstein statistics. The factor $(n_{\mathbf{k} \uparrow} + n_{-\mathbf{k} \downarrow})$ in (2-21b) represents the effect of the

Pauli principle acting on the individual electrons forming the pair. Perhaps it is simplest to view the "pairon" operators b_k and b_k^+ as satisfying Bose-Einstein statistics for $k \neq k'$ and satisfying the Pauli principle $b_k^{+2} = 0 = b_k^2$ for $k = k'$. It is the latter relation which ruins a simple Bose gas picture.⁴⁷

In attempting to find a variational estimate of the ground-state energy and wave function of H_{red} , the author tried to adapt the intermediate coupling approximation of Tomonaga, familiar in the coupled meson-nucleon problem and the polaron problem.^{48, 49} In these problems one assumes that successive bosons (mesons or phonons) are emitted into the *same* orbital state (about the proton or the electron, respectively). The form of the orbital state φ and the weight A_N of the $N/2$ boson state are then determined by minimizing the system energy. Lee, Low, and Pines^{49a} simplified the procedure for the polaron by assuming what is equivalent to a parameterized form for the weights A_N . Their wave function, after a canonical transformation has been made to eliminate the electron's coordinate from the problem, is

$$|\psi_0\rangle \propto \prod_k e^{g_k(a_k^+ + a_{-k})} |0\rangle \quad (2-22)$$

where the a^+ 's are phonon creation operators. The function g_k is essentially the Fourier transform of the orbital wave function φ of the phonons surrounding the electron.

The application of this physical idea to superconductivity is complicated by several features. First, the "pairon" operators do not truly satisfy Bose statistics and, second, the number of electrons is a definite number N_0 in our system, rather than being a probability distribution $|A_N|^2$ of finite width about the value N_0 . The author tried to describe the ground state of H_{red} by

$$|\psi_0\rangle \propto \prod_k e^{g_k b_k^+} |0\rangle = \prod_k (1 + g_k b_k^+) |0\rangle \quad (2-23)$$

where we have used the fact that $b_k^{+2} = 0$ in expanding out the exponential. The normalization integral is easily seen to be

$$\langle \psi_0 | \psi_0 \rangle = \prod_k (1 + |g_k|^2) \quad (2-24)$$

so that

$$|\psi_0\rangle = \prod_k \frac{1 + g_k b_k^+}{(1 + |g_k|^2)^{1/2}} |0\rangle \quad (2-25)$$

is a properly normalized state. By expanding out the infinite product one sees that $|\psi_0\rangle$ has a nonvanishing amplitude for all even numbers of electrons, 0, 2, 4, However, by choosing g_k appropriately the mean number of particles described by $|\psi_0\rangle$ can be adjusted to be the required number N_0 . As in the grand canonical ensemble, one can show that the width of the distribution is of order $N_0^{1/2}$ so that particle number fluctuations cause no difficulty in a large system.

Since we want to minimize the ground-state energy subject to the constraint

$$\langle\psi_0|N_{op}|\psi_0\rangle \equiv \langle\psi_0|\sum_{ks} n_{ks}|\psi_0\rangle = N_0 \quad (2-26)$$

we use the Lagrange multiplier scheme and minimize

$$\delta W = \delta\langle\psi_0|H_{\text{red}} - \mu N_{op}|\psi_0\rangle = 0 \quad (2-27)$$

On combining (2-20) and (2-27) one finds the quantity to be minimized is⁸

$$W = \sum_k 2(\epsilon_k - \mu)v_k^2 + \sum_{k,k'} V_{k'k} u_k v_k u_{k'} v_{k'} \quad (2-28)$$

where u_k and v_k are defined by

$$u_k = \frac{1}{(1 + g_k^2)^{1/2}} \quad (2-29a)$$

and

$$v_k = \frac{g_k}{(1 + g_k^2)^{1/2}} \quad (2-29b)$$

thus

$$u_k^2 + v_k^2 = 1 \quad (2-29c)$$

We have assumed phases are chosen so that $V_{k'k}$ and g_k are real quantities. On minimizing W one finds

$$u_k^2 = \frac{1}{2} \left(1 + \frac{\epsilon_k - \mu}{E_k} \right) \quad (2-30a)$$

$$v_k^2 = \frac{1}{2} \left(1 - \frac{\epsilon_k - \mu}{E_k} \right) \quad (2-30b)$$

and

$$u_k v_k = \frac{\Delta_k}{2E_k} \quad (2-30c)$$

where E_k is defined by

$$E_k = +[(\epsilon_k - \mu)^2 + \Delta_k^2]^{1/2} \quad (2-30d)$$

As we shall see, E_k turns out to be the energy required to create a quasi-particle of momentum \mathbf{k} in the superconducting state. The "energy-gap" parameter Δ_k satisfies the integral equation

$$\Delta_k = - \sum_{k'} V_{kk'} \frac{\Delta_{k'}}{2E_{k'}} \quad (2-30e)$$

One must simultaneously solve (2-30e) and the constraint condition

$$\langle \psi_0 | N_{op} | \psi_0 \rangle = 2 \sum_{\mathbf{k}} v_{\mathbf{k}}^2 = N_0 \quad (2-31)$$

to determine Δ_k and μ . If the single-particle energy ϵ_k is measured relative to the Fermi energy in the normal state, μ is just the shift of the chemical potential between the normal and superconducting states. For a system possessing particle-hole symmetry in the vicinity of the Fermi surface, one finds $\mu = 0$. In general, this is an excellent approximation and we shall assume $\mu = 0$ in solving for Δ_k . Once the energy-gap equation (2-30e) is solved, one can obtain the energy difference $W_N - W_S$ between the N - and S -states by inserting the expressions for u_k and v_k back into (2-28). An explicit solution of (2-30e) is easily obtained if $V_{kk'}$ is approximated by the s -wave potential

$$V_{kk'} \equiv \begin{cases} -V < 0 & \text{for } |\epsilon_k| \text{ and } |\epsilon_{k'}| < \omega_c \\ 0 & \text{otherwise} \end{cases} \quad (2-32)$$

so that $V_{kk'}$ is attractive in a shell of width $2\omega_c$ centered at the Fermi surface. In this case one finds

$$\Delta_k = \begin{cases} \Delta_0 & \text{for } |\epsilon_k| < \omega_c \\ 0 & \text{otherwise} \end{cases} \quad (2-33)$$

where

$$\Delta_0 = \frac{\omega_c}{\sinh \left[\frac{1}{N(0)V} \right]} \simeq 2\omega_c \exp \left[-\frac{1}{N(0)V} \right] \quad (2-34)$$

42 Theory of Superconductivity

The right-hand equality holds in the weak-coupling limit $N(0)V \gtrsim \frac{1}{4}$. By substituting this result into the expression for the ground-state energy (2-28) and subtracting the ground-state energy of the normal phase (i.e., the unperturbed Fermi sea in this model) one finds the condensation energy

$$W_N - W_S = \frac{1}{2} N(0) \Delta_0^2 \cong 2N(0)\omega_c^2 \exp \left[-\frac{2}{N(0)V} \right] \quad (2-35)$$

Since thermodynamics⁴ gives the relation

$$W_N - W_S = \frac{H_0^2}{8\pi} \quad (2-36)$$

where H_0 is the critical magnetic field for destroying superconductivity at zero temperature, we find

$$H_0 = 2[\pi N(0)]^{1/2} \Delta_0 \quad (2-37)$$

By using experimental values of $N(0)$ and Δ_0 one finds values of H_0 which are in reasonably good agreement with experiment.^{9, 16}

We note that the condensation energy (2-35) is not an analytic function of the coupling constant $N(0)V$ so that a perturbation treatment starting from the normal phase could not give this result unless one sums an infinite number of graphs of a selected class.⁵⁰

Returning to the ground-state wave function (2-23), we would expect on the basis of the Tomonaga scheme that the projection of $|\psi_0\rangle$ onto the N -particle space would lead to a function (in the coordinate representation) of the form

$$\begin{aligned} \langle r_1, s_1; r_2, s_2; \cdots r_N, s_N | \psi_0 \rangle &\equiv \psi_{0N} \\ &= \mathcal{A} \varphi(\mathbf{r}_1 - \mathbf{r}_2) \chi_{12} \varphi(\mathbf{r}_3 - \mathbf{r}_4) \chi_{34} \cdots \varphi(\mathbf{r}_{N-1} - \mathbf{r}_N) \chi_{N-1, N} \end{aligned} \quad (2-38)$$

The function φ is the relative coordinate wave function of a pair (*the same function for all pairs*) and χ_{ij} is corresponding spin function $\uparrow(i)\downarrow(j)$. Thus within the pairing approximation all pairs are in the *same* state in the ground-state wave function. The operator \mathcal{A} in (2-38) antisymmetrizes the entire function.

This result, first noted by Dyson, can be obtained by taking the inner product of $|\psi_0\rangle$ with the basis vector

$$|\mathbf{r}_1, s_1; \mathbf{r}_2, s_2; \dots; \mathbf{r}_N, s_N\rangle = \psi_{s_1}^+(\mathbf{r}_1)\psi_{s_2}^+(\mathbf{r}_2)\dots\psi_{s_N}^+(\mathbf{r}_N)|0\rangle$$

and expanding the ψ^+ 's in terms of the creation operators $c_{ks_i}^+$ by

$$\psi_{s_i}^+(\mathbf{r}_i) = \sum_k e^{-i\mathbf{k}\cdot\mathbf{r}_i} c_{ks_i}^+ \quad (2-39)$$

The N -particle state (2-38) has been discussed by Blatt^{51a} in the case where φ is a general function of \mathbf{r}_1 and \mathbf{r}_2 . This generalization corresponds to considering pairings between states other than $\mathbf{k}\uparrow$ and $-\mathbf{k}\downarrow$.

The orbital function φ in (2-38) is given by

$$\varphi(\boldsymbol{\rho}) = \sum_k g_k e^{i\mathbf{k}\cdot\boldsymbol{\rho}} \quad (2-40)$$

thus g_k is the Fourier transform of $\varphi(\boldsymbol{\rho})$, as stated above. While there is a formal similarity between (2-38) and the wave function for a condensed Bose-Einstein gas of pairs of electrons with opposite spin, the antisymmetrization operator \mathcal{A} is all important in real superconductors. In fact, the (unnormalized) ground state of the noninteracting Fermi gas can be written in this form with

$$g_k = \begin{cases} 1 & |k| < k_f \\ 0 & |k| > k_f \end{cases} \quad (2-41)$$

so that antisymmetrization removes the correlations between opposite spin electrons implied by φ in this case. In the superconducting state g_k differs from (2-41) only for values of \mathbf{k} in the immediate vicinity of the Fermi surface. This difference is reflected in a long-range tail of $\varphi(\boldsymbol{\rho})$ which increases the probability of two antiparallel spin electrons being near each other out to a range $\xi_0 \equiv \hbar v_F / \pi \Delta_0 \sim 10^{-4}$ cm, that is, Pippard's coherence length. As we mentioned above, this quasi-bound state has such a long range in space that on the average about 10^6 other pairs have their centers of mass in this region. [In this estimate electrons deep within the Fermi sea have not been counted since they behave essentially as if the material were in the normal phase.] Thus an isolated pair picture has little meaning here.