

# “Tight Binding” Method: Linear Combination of Atomic Orbitals (LCAO)

W. E. Pickett  
(Dated: April 14, 2014)

This write-up is a homemade introduction to the tight binding representation of the electronic structure of crystalline solids. This information is important in the parametrization of the band structures of real solids and for the underlying character of model Hamiltonians for correlated electron studies, as well as for other uses. Those interested in the parametrization of band structures of real materials should consult the book by

D. A. Papaconstantopoulos,  
*Handbook of the Band Structure of Elemental Solids*, (Plenum, New York, 1986).

For the general TB method, one should consult the original classic work by

J. C. Slater and G. F. Koster, Phys. Rev. **94**, 1498 (1954).

## I. INTRODUCTION TO TIGHT BINDING THEORY

Since a crystal is made up of a periodic array of atoms, it may seem peculiar that when we think of Bloch electron wavefunctions in solids it is often in terms of wavy modulations that don't pay much attention to just where the atoms sit. Indeed, in simple metals and covalent semiconductors that is a good picture: the crystal potential that enters into the Hamiltonian is a smooth function and atomic sites *per se* are not critical in the understanding (although they are in the underlying description of covalent semiconductors).

There is in fact a common picture – the tight binding model – that is based on the “collection of atoms” viewpoint. It is most appropriate when electrons move through the crystal slowly (or not at all, as in insulators) and therefore ‘belong’ to an atom for an appreciable time before they move on. The electrons are in some sense *tightly bound* to the atom and only hop because staying put on a simple atom costs a bit too much energy. The TB model is not readily applicable to simple (free or nearly free electron) metals, but it is quite good for a wide variety of other solids. It is interesting that covalent semiconductors can be described well from either viewpoint.

### A. Crystal as a collection of atoms

A good approximation for the electron's potential  $V(\vec{r})$  in a crystal is the sum of atomic potentials:

$$V(\vec{r}) = \sum_{\vec{R}} V_{at}(\vec{r} - \vec{R}), \quad (1.1)$$

where the sum runs over lattice vectors. We will not worry about the considerable non-uniqueness of this decomposition. For parametrization purposes, this

non-uniqueness is irrelevant (only the *form* of the resulting parameters is relevant), while if the method is intended for real electronic structure calculations, the non-uniqueness often is used to make numerical procedures as convenient as possible.

This potential is periodic by construction:

$$\begin{aligned} V(\vec{r} + \vec{R}_o) &= \sum_{\vec{R}} V_{at}(\vec{r} + \vec{R}_o - \vec{R}) \\ &= \sum_{\vec{R}} V_{at}(\vec{r} - (\vec{R} - \vec{R}_o)) \\ &= \sum_{\vec{R}'} V_{at}(\vec{r} - \vec{R}') \\ &\equiv V(\vec{r}), \end{aligned} \quad (1.2)$$

where the change of summation index  $\vec{R} \rightarrow \vec{R}' = \vec{R} - \vec{R}_o$  was made. Here  $\vec{R}_o$  is any lattice vector.

The crystal Hamiltonian is, using units where  $\hbar^2/2m \equiv 1$ ,

$$H = -\nabla^2 + V(\vec{r}). \quad (1.3)$$

The potential  $V(\vec{r})$  is periodic so  $H$  is periodic. Later we will generalize the situation to cover several atoms in the unit cell.

### B. Periodic array of atomic orbitals

Shouldn't the electron wavefunction in the crystal be related to the *atomic* orbitals, which satisfy

$$H_{at}\phi_n \equiv (-\nabla^2 + V_{at})\phi_n = \varepsilon_n\phi_n. \quad (1.4)$$

We might try the most simple linear combination of atomic orbitals that is periodic

$$\Phi_n(\vec{r}) = \sum_{\vec{R}} \phi_n(\vec{r} - \vec{R}); \quad (1.5)$$

is this a Bloch function that can be put into the form

$$\Phi(\vec{r}) = e^{i\vec{k}\cdot\vec{r}} u_{\vec{k}}(\vec{r})? \quad (1.6)$$

Indeed it is, *but* only for  $\vec{k} = 0$ . We want, and need, Bloch-like functions for arbitrary  $\vec{k}$  within the first Brillouin zone (BZ).

**Problem 1: prove this statement, that this proposed function has Bloch form for, and only for,  $\vec{k}=0$ .**

### C. Bloch Sums

A much better choice of candidate for a crystal wavefunction is to form the “Bloch sums” of atomic orbitals, given by

$$B_{n,\vec{k}}(\vec{r}) = N^{-\frac{1}{2}} \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} \phi_n(\vec{r} - \vec{R}). \quad (1.7)$$

where  $\vec{R}$  sums over the  $N$  lattice sites in our normalization volume. This type of linear combination is called a Bloch sum because it produces a function that satisfies the Bloch condition for wavevector  $\vec{k}$ :

$$\begin{aligned} N^{\frac{1}{2}} B_{n,\vec{k}}(\vec{r} + \vec{R}_o) &= \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} \phi_n(\vec{r} + \vec{R}_o - \vec{R}) \\ &= \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} \phi_n(\vec{r} - (\vec{R} - \vec{R}_o)) \\ &= \sum_{\vec{R}'} e^{i\vec{k}\cdot(\vec{R}' + \vec{R}_o)} \phi_n(\vec{r} - \vec{R}') \\ &= e^{i\vec{k}\cdot\vec{R}_o} \sum_{\vec{R}'} e^{i\vec{k}\cdot\vec{R}'} \phi_n(\vec{r} - \vec{R}'), \\ &= N^{\frac{1}{2}} e^{i\vec{k}\cdot\vec{R}_o} B_{n,\vec{k}}(\vec{r}). \end{aligned} \quad (1.8)$$

so

$$B_{n,\vec{k}}(\vec{r} + \vec{R}_o) = e^{i\vec{k}\cdot\vec{R}_o} B_{n,\vec{k}}(\vec{r}). \quad (1.9)$$

This result then is in Bloch form. All such Bloch functions are periodic in  $\vec{k}$  from BZ to BZ.

**Problem 2: prove this statement by manipulating it into the Bloch form, and discover what periodic part  $u_{\vec{k}}(\vec{r})$  is.**

It suffices to confine  $\vec{k}$  in Eq. 1.7 to the 1st BZ. If the  $\vec{k}$  point were of the form  $\vec{k}_r + \vec{K}$ , where  $\vec{k}_r$  (the *reduced wavevector*) is in the 1st BZ and  $\vec{K}$  is a reciprocal lattice vector, note that

$$e^{i\vec{k}\cdot\vec{R}} = e^{i\vec{k}_r\cdot\vec{R}} \quad (1.10)$$

for any and all lattice vectors  $\vec{R}$ . (Be sure you understand why.) Thus these Bloch sums, and any functions constructed from them, are periodic from BZ to BZ.

### D. Proceeding toward the eigenfunctions <sup>2</sup>

For many solids there will be several types ( $s, p$  or  $d$ ) of atomic states in the valence region, and that is why we have kept the index  $n$ . In the solid these atomic states will mix with each other due to the overlap of atomic orbitals on neighboring atoms (as we will see). A Bloch sum of atomic orbitals itself is not an eigenfunction for the crystal. It is important to allow the valence wavefunction in the solid to be some of *each* of the atomic functions, with the actual amounts to be determined by solving Schrödinger's equation. Thus we try expressing the electron wavefunction in the crystal as a bit  $b_{n,\vec{k}}$  of each of the Bloch sums,

$$\psi_{\vec{k}}(\vec{r}) = \sum_n b_n(\vec{k}) B_{n,\vec{k}}(\vec{r}), \quad (1.11)$$

where the coefficients  $b$  gives the amount of Bloch sum  $B_n$  in the crystal wavefunction. The Bloch sums become the *basis functions* that we express the wavefunction in terms of. Since the Bloch sums themselves are normalized to unity, we will want  $\sum_n |b_n(\vec{k})|^2 = 1$  for each  $\vec{k}$ .

Now we want to learn how to find the wavefunctions. The condition is that they be solutions of the Schrödinger equation

$$H\psi_{\vec{k}} = \varepsilon_{\vec{k}} \psi_{\vec{k}}. \quad (1.12)$$

The eigenvalues  $\varepsilon_{\vec{k}}$  will be the *energy bands* of the crystal. But how do we solve for  $\varepsilon_{\vec{k}}$  and the coefficients  $b_n(\vec{k})$ ?

### E. The matrix equation

In quantum mechanics generally, a good try is to take *matrix elements* (integrals between basis functions) and reduce the problem to a matrix equation. In this case, the thing to do is to multiply the Schrödinger equation on the left by another Bloch sum  $B_{m,\vec{k}}^*$  and integrate over the crystal. Note that we have not chosen a Bloch function corresponding to another wavevector  $\vec{k}' \neq \vec{k}$ .

**Problem 3: calculate  $\langle B_{m\vec{k}} | B_{n\vec{k}'} \rangle$  to see why we simplify to  $k' = k$ . See below for notation.**

The result is

$$\sum_n H_{m,n}(\vec{k}) b_n(\vec{k}) = \varepsilon_{\vec{k}} \sum_n S_{m,n}(\vec{k}) b_n(\vec{k}), \quad (1.13)$$

where

$$\begin{aligned} H_{m,n}(\vec{k}) &\equiv \int B_{m,\vec{k}}^*(\vec{r}) H B_{n,\vec{k}}(\vec{r}) \\ &= \langle B_{m\vec{k}} | H | B_{n\vec{k}} \rangle \end{aligned} \quad (1.14)$$

$$S_{m,n}(\vec{k}) \equiv \int B_{m,\vec{k}}^*(\vec{r}) B_{n,\vec{k}}(\vec{r}) = \langle B_{m,k} | S_{n,k} | B_{n,k} \rangle$$

These matrices are called the Hamiltonian matrix and the overlap matrix, respectively, where  $m$  and  $n$  are the matrix indices. Writing the matrices implicitly (without displaying the indices), the equation becomes

$$H(\vec{k})b(\vec{k}) = \varepsilon_{\vec{k}} S(\vec{k})b(\vec{k}) \quad (1.16)$$

or

$$\{H(\vec{k}) - \varepsilon_{\vec{k}} S(\vec{k})\}b(\vec{k}) = 0. \quad (1.17)$$

This is a linear algebra problem (generalized eigenvalue problem), but we have to know what the matrices  $H$  and  $S$  are.

### F. The $H$ and $S$ matrices

Well, we never said that this wouldn't get a little messy. However, the messiness does clear up soon. Substituting in the Bloch sum forms for  $B_m^*$  and  $B_n$  within the integral, we have the expression

$$\begin{aligned} H_{m,n}(\vec{k}) &= (N^{-\frac{1}{2}})^2 \sum_{\vec{R}_1, \vec{R}_2} e^{i\vec{k} \cdot (\vec{R}_2 - \vec{R}_1)} \\ &\times \int \phi_m^*(\vec{r} - \vec{R}_1) H \phi_n(\vec{r} - \vec{R}_2) \\ &\equiv \frac{1}{N} \sum_{\vec{R}_1, \vec{R}_2} e^{i\vec{k} \cdot (\vec{R}_2 - \vec{R}_1)} H_{m,n}(\vec{R}_2 - \vec{R}_1). \end{aligned} \quad (1.18)$$

Because the Hamiltonian is cell periodic, the matrix element on the right hand side depends only on the difference between  $\vec{R}_2$  and  $\vec{R}_1$ . We can then change the summation index  $\vec{R}_2$  to  $\vec{R}_2 - \vec{R}_1 = \vec{R}$ , and the index  $\vec{R}_1$  no longer appears on the right hand side. Then the sum over  $\vec{R}_1$  just gives the factor  $N$ , the number of unit cells in our “crystal.” Then we obtain

$$H_{m,n}(\vec{k}) = \sum_{\vec{R}} e^{i\vec{k} \cdot \vec{R}} H_{m,n}(\vec{R}). \quad (1.19)$$

There should be no confusion between  $H_{m,n}(\vec{k})$  and  $H_{m,n}(\vec{R})$  in practice; in fact, they are *lattice Fourier transforms* of each other.

The expression for the real space integral is

$$H_{m,n}(\vec{R}) = \int \phi_m^*(\vec{r}) H \phi_n(\vec{r} - \vec{R}), \quad (1.20)$$

*i.e.* it indicates the amount by which the Hamiltonian  $H$  couples atomic orbital  $\phi_m$  on the site at the origin to the atomic orbital  $\phi_n$  that is located at site  $\vec{R}$ . Physically,  $H_{m,n}(\vec{R})$  is the amplitude that an electron in orbital  $\phi_n$  at site  $\vec{R}$  will hop to the orbital  $\phi_m$  at the origin under the action of the Hamiltonian. One limit is easy to see: if  $|\vec{R}|$  is so large that either one or the other of the orbitals is vanishingly small where the other is nonzero, then the integral is negligible. Thus we can confine ourselves to values of  $\vec{R}$  that connect an atom to only a few near neighbors. Often the “nearest neighbor” approximation is excellent.

All of this discussion applies as well to  $S$ , by just removing the Hamiltonian  $H$  from inside the integral.  $S_{m,n}(\vec{R})$  in fact is called the *overlap* of  $\phi_m(\vec{r})$  and  $\phi_n(\vec{r} - \vec{R})$ . Note that, if the orbitals are normalized (as is always possible, and is always assumed), the  $S_{m,m}(\vec{0})=1$  for all  $m$ , *i.e.* the diagonal elements of  $S$  are unity.

### H. Several atoms in the unit cell

So far, the notation has been limited to a elemental crystal. More generally, one encounters compounds where there are various atoms in the cell, which can be labeled by their position  $\vec{\tau}_i$  with respect to the origin of the cell ( $\vec{R}$ ). Then the basis orbitals are

$$\phi_{m,i,R} \equiv \phi_{m,i}(\vec{r} - \vec{R} - \vec{\tau}_i). \quad (1.21)$$

Then, generalizing from the Bloch sum defined in Eq. 1.7, one has the basis Bloch sums

$$B_{m,i,\vec{k}}(\vec{r}) = N^{-\frac{1}{2}} \sum_{\vec{R}} e^{i\vec{k} \cdot (\vec{R} + \vec{\tau}_i)} \phi_{m,i}(\vec{r} - \vec{R} - \vec{\tau}_i). \quad (1.22)$$

Then, instead of Eq. 1.18 we obtain

$$\begin{aligned} H_{m,i;n,j}(\vec{k}) &= (N^{-\frac{1}{2}})^2 \sum_{\vec{R}_1, \vec{R}_2} e^{i\vec{k} \cdot (\vec{R}_2 + \vec{\tau}_j - \vec{R}_1 - \vec{\tau}_i)} \\ &\times \int \phi_m^*(\vec{r} - \vec{R}_1 - \vec{\tau}_i) H \phi_n(\vec{r} - \vec{R}_2 - \vec{\tau}_j) \\ &= \frac{1}{N} e^{i\vec{k} \cdot (\vec{\tau}_j - \vec{\tau}_i)} \sum_{\vec{R}_1, \vec{R}_2} e^{i\vec{k} \cdot (\vec{R}_2 - \vec{R}_1)} \end{aligned}$$

$$\begin{aligned}
& \times H_{m,i;n,j}(\vec{R}_2 + \vec{\tau}_j - \vec{R}_1 - \vec{\tau}_i) \\
& = e^{-i\vec{k}\cdot\vec{\tau}_i} \left( \sum_{\vec{R}} H_{m,i;n,j}(\vec{R}) e^{i\vec{k}\cdot\vec{R}} \right) e^{i\vec{k}\cdot\vec{\tau}_j} \\
& = e^{-i\vec{k}\cdot\vec{\tau}_i} H_{m,i;n,j}^o(\vec{k}) e^{i\vec{k}\cdot\vec{\tau}_j}. \quad (1.23)
\end{aligned}$$

Here the notation in the next-to-last line has been shortened by leaving the internal coordinates implicit,

$$H_{m,i;n,j}(\vec{R}) \equiv H_{m,i;n,j}(\vec{R} + \vec{\tau}_j - \vec{\tau}_i). \quad (1.24)$$

The overlap matrix  $s$  behaves in a formally analogous way.

Eq. 1.23 can be viewed as the matrix  $\hat{H}^0(\vec{k})$  transformed by the unitary transformation

$$U_{m,i;n,j}(\vec{k}) = e^{i\vec{k}\cdot\vec{\tau}_j} \delta_{n,m} \delta_{i,j}, \quad (1.25)$$

which can easily be shown to obey the unitarity conditions

$$\hat{U}^\dagger \hat{U} = \hat{1} = \hat{U}^{-1} \hat{U}; \quad U^\dagger = \hat{U}^{-1}. \quad (1.26)$$

However, a unitary transformation of a Hermitian matrix does not affect its eigenvalues, but merely transforms the eigenvectors. So, unless there is some specific reason for doing so, the phase factors in the last line of Eq. 1.23) (*i.e.*  $\hat{U}$ ) can be disregarded. Looked at the other way, the one may *include* any additional unitary transformation that one desires. There are occasionally good reasons for making the transformation – for checking various aspects of the code, for example, or more importantly for reducing the secular equation to a real equation in cases where there is a center of inversion in the crystal but the original choice of orbitals produced a complex secular equation.

The important result is that, for obtaining the eigenenergies (the energy bands) and even fractions of atomic orbitals of each eigenstate, one does not need to deal with the internal atomic positions in constructing  $H_{m,i;n,j}(\vec{k})$  and this simplification is usually made.

## II. APPLICATIONS OF THE TIGHT BINDING MODEL

It may not have been obvious, but the mathematics is finished; the problem has been solved except for the numerical work. Obtaining the energy bands  $\varepsilon_{\vec{k}}$  (in general there will be several of them, labeled  $\varepsilon_{\vec{k},n}$ ) and the expansion coefficients  $b_n(\vec{k})$  of the wavefunctions amounts to solving Eq. 1.16 or 1.17, with the matrix  $H$  given by Eq. 1.19 and  $S$

given by an analogous expression:

$$\begin{aligned}
S_{m,n}(\vec{k}) &= N^{-1} \sum_{\vec{R}_1, \vec{R}_2} e^{i\vec{k}\cdot(\vec{R}_2 - \vec{R}_1)} S_{m,n}(\vec{R}_2 - \vec{R}_1) \\
&= \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} S_{m,n}(\vec{R}). \quad (2.27)
\end{aligned}$$

### A. Single Site Terms

First we will look at the  $\vec{R} = \vec{0}$  terms, where both orbitals are on the same site (at our origin). Now it is helpful to write the crystal Hamiltonian as the atomic Hamiltonian for the atom at the origin, plus the potential from all of the other atoms:

$$\begin{aligned}
H &= -\nabla^2 + V_{at}(\vec{r}) + \sum_{\vec{R} \neq 0} V_{at}(\vec{r} - \vec{R}) \\
&= -\nabla^2 + V_{at}^{sph}(|\vec{r}|) + V_{at}^{non}(\vec{r}) + \sum_{\vec{R} \neq 0} V_{at}(\vec{r} - \vec{R}) \\
&= H_{at}^{sph}(\vec{r}) + \Delta V(\vec{r}). \quad (2.28)
\end{aligned}$$

The integral results primarily from the first part, which gives the “atomic” eigenvalues  $\{\varepsilon_m\}$  for a spherical atomic Hamiltonian  $H_{at}^{sph}$ . Then because atomic orbitals on the same atom are orthogonal to each other, the on-site matrix element is

$$\int \phi_m^*(\vec{r}) H_{at}(\vec{r}) \phi_n(\vec{r}) = \varepsilon_n \delta_{m,n}, \quad (2.29)$$

*i.e.* just the atomic eigenvalue.

### B. Crystal Field Splitting

The quantity  $\Delta V(\vec{r})$  in Eq. 2.28 is not well specified, but we do know that it has the symmetry of the atom in questions (defining the origin in this equation). This symmetry is never spherical in a solid, but discrete, such as a mirror plane ( $m$ ), a 3-fold rotation in an axis containing inversion together with a mirror plane ( $\bar{3}m$ ), etc. This *crystal field*, that is, the non-spherical potential that arises because the atom is in a crystal, will split some eigenvalues that would be degenerate in a spherical potential. The most common such situation is for the five  $d$  orbitals in a cubic crystal field, which are split into the threefold representation called  $t_{2g}$  ( $xy, yz, zx$ ) and the twofold representation known as  $e_g$  ( $x^2 - y^2, 3z^2 - 1$ ). Crystal fields are also very important for  $4f$  and  $5f$  ions, where the designation may become more complex because intra-atomic coupling (correlation) is important. Although not widely acknowledged, the oxygen ion in an axial site (such as in the perovskite

structure) has crystal field split  $p$  levels: twofold  $(x, y)$  and a singlet  $(z)$ . What this means is that instead of a single on-site (“atomic”) energy  $\varepsilon_d$  for a transition metal ion in a cubic site, one has two energies  $\varepsilon_{t_{2g}}$  and  $\varepsilon_{e_g}$ , or even more distinct ones if the symmetry is lower.

### C. Three Center vs. Two Center

A general Hamiltonian matrix element in Eq. (1.20) contains atomic potentials on a third atom, besides the sites upon which the orbitals are centered. The general form of the matrix elements then contains *three center integrals*. Slater and Koster introduced, and advocated, the use of the *two-center approximation* (2CA), in which three center contributions to the parameters are neglected. By the way, if you can follow this set of notes to this point, you can read and understand the Slater-Koster that is referenced in the abstract. It is a must-read for those using the tight binding method.

There had already been suggestions that such three center terms were negligible and, while pointing out that they are not negligible in a serious calculation (borne out by many subsequent studies), SK suggested that for the purposes of parametrizing information (from experiment or from band calculation) a 2CA is likely to be very useful. Table I in their paper provides the expression for  $s, p$ , and  $d$  matrix elements in terms of *two center integrals*, denoted  $(ss\sigma), (pd\pi), (dd\delta)$ , etc. The 2CA is very widely, almost universally, used.

As an example of the validity, or lack thereof, of the 2CA, we consider the case of Nb studied by Pickett and Allen.[1] They performed a full three center fit to the augmented plane wave bands of Mattheiss for Nb (also Mo; these were state of the art in accuracy at the time, and still perfectly good for most applications) at 55  $k$  points in the irreducible zone, using an  $s + p + d$  basis set ( $9 \times 9$  matrix), and including parameters out to third neighbors. Several of the 31 parameters are found to be negligibly small, suggesting the full three-center treatment was not essential. As a test of the 2CA in Nb, we can look at the  $pd$  parameters for nearest neighbors in the bcc lattice, which lie along (111) directions. The values of the independent parameters (in mRy), together with the expression in the 2CA, are

$$\begin{aligned} -42 &= H_{x,xy} \rightarrow \frac{1}{3}(pd\sigma) + \frac{1}{3\sqrt{3}}(pd\pi) \\ +16 &= H_{x,x^2-y^2} \rightarrow 0(pd\sigma) + \frac{1}{3\sqrt{3}}(pd\pi) \\ -51 &= H_{x,yz} \rightarrow \frac{1}{3}(pd\sigma) - \frac{2}{3\sqrt{3}}(pd\pi). \end{aligned}$$

Since the 2nd equation determines  $(pd\pi) = 28 \text{ mRy}$  ( $= 0.38 \text{ eV}$ ), the other two equations can be used separately to “determine”  $(pd\sigma)$ . The two values thus determined are -142 mRy and -131 mRy. Since these do not differ greatly, one can use the 2CA with  $(pd\sigma) = -136 \text{ mRy}$  (the mean) with acceptable loss of accuracy.

The  $dd$  parameters are the crucial ones in Nb and in other elemental transition metals, and in most transition metal compounds. For nearest neighbors there are four three center parameters. Denoting  $(dd\sigma), (dd\pi), (dd\delta)$  by  $S, P, D$ , respectively, the corresponding values and 2CA expressions are given by

$$\begin{aligned} -22 &= H_{xy,xy} \rightarrow \frac{1}{3}S + 0P + \frac{4}{9}D \\ -35 &= H_{xy,zx} \rightarrow \frac{1}{3}S - \frac{1}{9}P - \frac{2}{9}D \\ -31 &= \sqrt{3}H_{xy,3z^2-r^2} \rightarrow 0S - \frac{2}{3}P + \frac{2}{3}D \\ +31 &= H_{3z^2-r^2,3z^2-r^2} \rightarrow \frac{2}{9}S + \frac{2}{3}P + \frac{1}{3}D \end{aligned} \quad (2.30)$$

The 3rd of these equations has been multiplied through by  $\sqrt{3}$  to facilitate solution by elimination using various combinations of the equations. Subtracting the 2nd from the 1st gives

$$+13 = 0S + \frac{1}{9}P + \frac{2}{3}D, \quad (2.31)$$

which can be combined with the 3rd and 4th equations (separately) to produce three “solutions,” one of which is  $\{S, P, D\} = \{-107, 77, 31\}$ . Notice the strong decrease from  $\sigma \rightarrow \pi \rightarrow \delta$ . Comparison of the values from different “solutions” provides a guideline for the viability of the 2CA.

### D. Introducing the usual terminology

Now we consider an intersite term. However, we do not intend here to do a serious calculation; rather, we want to identify the important quantities (often called *parameters*) and choose likely values and learn some things about the general behavior of the electronic system. In fact, we’ll just define a TB parameter

$$t_{m,n}(\vec{R}) \equiv H_{m,n}(\vec{R}) \quad (2.32)$$

because  $t$  is the usual notation for these integrals. This integral (see above) indicates how easy it is for an electron – whose behavior after all is determined by the Hamiltonian – to “hop” from orbital  $n$  on site  $\vec{R}$  to orbital  $m$  at the origin. The  $\{t\}$  constants are called *hopping parameters* or *hopping amplitudes*.

The on-site ( $\vec{R} = 0$ ) term was given in Eq. 2.29, which we repeat here to jog the memory:

$$t_{m,n}(\vec{0}) = \varepsilon_n \delta_{m,n}, \quad (2.33)$$

which is the atomic energy level corresponding to orbital  $\phi_n$ .

For the overlap matrix we have similarly the notation

$$s_{m,n}(\vec{R}) = S_{m,n}(\vec{R}); \quad s_{m,n}(0) = \delta_{m,n}. \quad (2.34)$$

The latter result expresses the orthonormality of atomic orbitals.

Note that the local orbitals have never been constructed. They enter implicitly through the Hamiltonian and overlap parameters; the method is a parametrization scheme. It is possible to do actual calculations using atomic-like orbitals, which is known as linear combination of atomic orbitals (LCAO).

### III. EXAMPLES

#### A. $s$ functions on a simple lattice

The simplest case to consider is when there is only one,  $s$ -like function on each atom at sites in a Bravais lattice (*i.e.* one atom per cell). Then the TB matrix equation is a  $1 \times 1$  equation, which gives a direct expression for  $\varepsilon_{\vec{k}}$ . Since an  $s$  atomic orbital is a spherically symmetric function, there is no aggravating angular dependence to worry about in Eq. ?? for the hopping parameters, and in fact the value of the hopping parameter is the same for all equivalent neighbors – all 1st neighbors have the same value, all 2nd neighbors have the same (different, usually smaller) value, etc. Let's look at some cases.

#### B. 1D linear chain of atoms

Taking some atom as our origin, we have two 1st neighbors, at  $\pm a$ , and both have the same hopping amplitude  $t_1$ . Then the sum in Eq. 1.19 contains the on-site term and those from neighbors:

$$\begin{aligned} H_{s,s}(k) &= \varepsilon_s + t_1 \sum_{\vec{R}} e^{ikR} \\ &= \varepsilon_s + t_1 (e^{ikR} + e^{-ikR}) \\ &= \varepsilon_s + 2t_1 \cos(ka). \end{aligned} \quad (3.35)$$

Likewise,

$$S_{s,s}(k) = 1 + s_1 \sum_{\vec{R}} e^{ikR} = 1 + 2s_1 \cos(ka). \quad (3.36)$$

The solution to Eq. 1.16 is immediate:

$$\varepsilon_k = \frac{\varepsilon_s + 2t_1 \cos(ka)}{1 + 2s_1 \cos(ka)} \quad (3.37)$$

It is very simple to add the effects of interaction with 2nd neighbors, which lie at  $\pm 2a$  from the reference atom (“at the origin”). The sum over the complex exponential factors leads to the result  $2\cos(2ka)$ . Denoting the corresponding integrals by  $t_2$  and  $s_2$ , the energy band *dispersion relation* becomes

$$\varepsilon_k = \frac{\varepsilon_s + 2t_1 \cos(ka) + 2t_2 \cos(2ka)}{1 + 2s_1 \cos(ka) + 2s_2 \cos(2ka)} \quad (3.38)$$

#### C. General Features

It is time to reflect. Eq. 3.35 is the simple *1D tight binding band* that arises commonly in modeling the behavior of 1D materials. Actually, it is almost always even simpler, because the  $s$  overlap parameters are usually neglected for further simplicity. This is discussed in the next subsection. However, at the cost of a little messiness in deriving the expression, we have obtained a very simple form of  $\varepsilon$  vs.  $k$  dispersion relation  $\varepsilon_k$ . In the simplest case it has only a single parameter,  $t_1 \equiv t$ , and by construction – because the wavefunctions were built to satisfy the Bloch condition – it has precisely the correct periodicity. These are two general features of the TB representation: (i) simplicity, and (ii) expressions involving trig functions that automatically have the correct periodicity.

Another physical requirement is that the parameter  $S_{n,m}(\vec{R})$  in Eq. 2.27 cannot be greater than unity in absolute value, however there can be a problem with the denominator vanishing in Eq. 3.37 unless  $s_1$  is less than 0.5. In practice, to be reasonable the various overlap parameters  $s_{m,n}(\vec{R})$  should be small in magnitude compared with unity (they may be of either sign). The “self overlap” term is unity by normalization, and putting one of the orbitals on a different site must reduce the magnitude of the overlap. Hence the denominator in Eq. 3.37 should always take the form of a correction, an *adjustment*, and not give very large alteration of the denominator, or else the description loses its realism.

If the overlap matrix is approximated by the unit matrix, so that Eq. 3.37 becomes

$$\varepsilon_k = \varepsilon_s + 2t_1 \cos(ka), \quad (3.39)$$

the interpretation of the parameters occurring in the dispersion relation is simple. The band minimum and maximum are  $\varepsilon_s - |2t_1|$  and  $\varepsilon_s + |2t_1|$  respectively. This means that the center of the band lies

$$W = 4|t_1|. \quad (3.40)$$

This generalizes to square/cubic lattices in 2D and 3D, where the bandwidth is given by  $2z|t_1|$ , where  $z$  is the usual notation for the number of nearest neighbors. This can be changed somewhat by 2nd neighbor terms, but the rule-of-thumb is this simple relationship between bandwidth, coordination number, and nearest neighbor hopping amplitude:

$$W = 2z|t_1|. \quad (3.41)$$

#### D. Character of the Overlap Correction

We can obtain insight into the effect of including the overlap matrix  $S$  by expanding Eq. 3.38, assuming  $s_1, s_2$  are small. We obtain, shortening  $\cos(ka) \rightarrow C_1(k), \cos(2ka) \rightarrow C_2(k)$ ,

$$\begin{aligned} \varepsilon_k &\approx (\varepsilon_s + 2t_1C_1(k) + 2t_2C_2(k)) \times \\ &\quad (1 - 2s_1C_1(k) - 2s_2C_2(k)) \\ &= \varepsilon_s + 2(t_1 - s_1)C_1(k) + 2(t_2 - s_2)C_2(k) \\ &\quad - 4t_1s_1C_1(k)^2 - 4t_2s_2C_2(k)^2 \\ &\quad - 4(t_1s_2 + t_2s_1)C_1(k)C_2(k). \end{aligned} \quad (3.42)$$

Several effects can be noted. In the simple  $C_1(k)$  and  $C_2(k)$  terms, the overlaps are reduced from  $t_1, t_2$  to  $t_1 - s_1$  and  $t_2 - s_2$ . Of course this can be an increase if  $t_1$  and  $s_1$  differ in sign, etc. The second effect is to add more wiggles (Fourier components), via the products of  $c_j(k)$  terms. It does so in a somewhat different way than simply adding more neighbors (more  $t$ 's) into the expansion. Probably it is more efficient, when using the TB parametrization to fit given complicated band structures, to use an overlap matrix rather than simply add more hopping parameters, but this question may not have been studied systematically.

#### IV. THE OVERLAP MATRIX

The overlap matrix deserves comment before we continue. In a large majority of cases where the TB method is used for simplification or for pedagogical reasons, the overlap matrix is “set to unity”:

$$S_{m,n}(\vec{R}) \rightarrow \delta_{m,n} \delta_{\vec{R}, \vec{0}}. \quad (4.43)$$

It is important to understand why, first, this is in principle a reasonable thing to do, and second, it leads not only to simplification but to additional approximation.

Let us return to Eq 1.17,

$$\hat{H}b = \varepsilon \hat{S}b, \quad (4.44)$$

where the notation has been simplified even further by dropping the  $\vec{k}$  argument on each quantity, but adding a “hat” on the matrices. We can make a transformation that effectively eliminates  $S$ .

This is done by introducing the square root, denote it  $\hat{S}^{\frac{1}{2}}$ , of the overlap matrix  $S$ :

$$\hat{S} = \hat{S}^{\frac{1}{2}} \hat{S}^{\frac{1}{2}} = (\hat{S}^{\frac{1}{2}})^2. \quad (4.45)$$

We will also need the *inverse* of the square root of  $\hat{S}$  (equal to the square root of the inverse):

$$\hat{S}^{-\frac{1}{2}} \hat{S}^{\frac{1}{2}} = \hat{1} = \hat{S}^{\frac{1}{2}} \hat{S}^{-\frac{1}{2}}. \quad (4.46)$$

Aha! I hear you say. Square roots and inverses of matrices do not always exist. That is true; however,  $\hat{S}$  is a *positive matrix* from its physical and mathematical definition. This means that all of its eigenvalues are positive, in which case its square root and its inverse do exist. These matrices can be obtained by (1) performing the similarity transformation that transforms  $\hat{S}$  to a diagonal matrix, (2) taking the square root[2] or the inverse of all diagonal elements as desired, and (3) performing the inverse similarity transformation. It is readily shown that the corresponding matrices obey the properties given above for the square root or the inverse. We don't say more about this because for now it is only necessary to know that it is possible.

With these matrices we can now carry out the following steps:

$$\begin{aligned} \hat{H}b &= \varepsilon \hat{S}b \\ \hat{H} \hat{S}^{-\frac{1}{2}} \hat{S}^{\frac{1}{2}} b &= \varepsilon \hat{S}^{\frac{1}{2}} \hat{S}^{\frac{1}{2}} b \\ (\hat{S}^{-\frac{1}{2}} \hat{H} \hat{S}^{-\frac{1}{2}})(\hat{S}^{\frac{1}{2}} b) &= \varepsilon (\hat{S}^{\frac{1}{2}} b) \\ \hat{\mathcal{H}} \bar{b} &= \varepsilon \bar{b}, \end{aligned} \quad (4.47)$$

where

$$\hat{\mathcal{H}} = \hat{S}^{-\frac{1}{2}} \hat{H} \hat{S}^{-\frac{1}{2}}. \quad (4.48)$$

The generalized eigenvalue problem  $\hat{H}b = \varepsilon \hat{S}b$  has been transformed into a conventional eigenvalue problem  $\hat{\mathcal{H}}\bar{b} = \varepsilon \bar{b}$  with the same eigenvalues. The eigenvectors, which contain the information about how much each atomic orbital contributes to the eigenfunction, have changed, and the Hamiltonian matrix that needs to be diagonalized has been transformed. This observation, together with the lack of any overlap matrix in the new equation, indicates that the transformation

$$\bar{b} = \hat{S}^{\frac{1}{2}} b \quad (4.49)$$

reflects a transformation of the underlying atomic orbitals into a set of mutually orthogonal orbitals, *even if they were originally residing on different atoms*. This orthogonalization procedure is called Löwdin orthogonalization, after the eminent quantum chemist Per-Olov Löwdin (pronounced *love-dean*).

What this orthogonalization procedure does is to add in (either sign is possible) to any given orbital  $\phi_j(\vec{r})$  a fraction of every other orbital that it overlaps. The amount it must mix in is related to the corresponding element of  $S_{m,n}(\vec{R})$ . The effect is to make each of the orbitals more spread out, and thereby to increase the number of matrix elements that are needed in  $\hat{\mathcal{H}}$  compared to what were needed in  $\hat{H}$ . This fact is bothersome and tends to get “forgotten,” because the objective of the TB formalism is simplicity, and therefore few parameters. The conventional applications of TB theory for (semi)empirical studies therefore usually presumes (“we assume, for simplicity”) that the original local orbitals  $\{\phi_m(\vec{r} - \vec{R})\}$  have been orthogonalized.

If the off-diagonal elements of  $S$  (*i.e.* the overlaps) are small, an approximate square-root matrix can be obtained by expansion. Writing

$$\delta\hat{S} \equiv \hat{S} - 1, \quad (4.50)$$

we can write

$$\hat{S}^{\frac{1}{2}} = (1 + \delta\hat{S})^{1/2} \approx 1 + \frac{1}{2}\delta\hat{S} + \frac{1}{8}(\delta\hat{S})^2 + \dots \quad (4.51)$$

This expansion is readily evaluated without diagonalizing the overlap matrix.

## V. RETURN TO APPLICATIONS

### A. 2D Square Lattice of $s$ Orbitals

Henceforward we neglect the  $\hat{S}$  matrix, since most of the rest of the world does so. For the square lattice, the sum over nearest neighbors runs over the sites  $(a, 0), (0, a), (-a, 0), (0, -a)$ . The sum becomes

$$\begin{aligned} \sum_{\vec{R}} &\rightarrow \sum_{j=\pm 1} e^{ik_x a j} + \sum_{p=\pm 1} e^{ik_y a p} \\ &= 2(\cos(k_x a) + \cos(k_y a)), \end{aligned} \quad (5.52)$$

and the dispersion relation is

$$\varepsilon_{\vec{k}} = \varepsilon_s + 2t_1(\cos(k_x a) + \cos(k_y a)). \quad (5.53)$$

Now, suppose we want to account for “hopping” to the second neighbors at the points  $(\pm a, \pm a)$  with

amplitude  $t_2$ , we take advantage of  $e^{v+w} = e^v e^w$  to get the additional term, and the band looks like

$$\begin{aligned} \varepsilon_{\vec{k}} = \varepsilon_s &+ 2t_1(\cos(k_x a) + \cos(k_y a)) \\ &+ 4\cos(k_x a)\cos(k_y a). \end{aligned} \quad (5.54)$$

Once you do a few of these, the pattern becomes simple. When you need to deal with atomic  $p$  or  $d$  orbitals, the procedure becomes more intricate: the  $t(\vec{R})$  factor depends on  $\vec{R}$  and cannot simply be pulled out from under the summation. But that is another course.

### B. 2D square lattice of $p$ orbitals

For  $p_x, p_y$  orbitals on a square lattice, one can take advantage of the observation that  $p_x$  and  $p_y$  orbitals on neighboring sites are orthogonal, whether oriented along the  $\hat{x}$  or  $\hat{y}$  directions. Keeping the larger  $p\sigma$  hopping and also the smaller  $p\pi$  hopping, the  $2 \times 2$  matrix (for the two orbitals) is diagonal, with

$$p_x = 2t_\sigma \cos k_x a + 2t_\pi \cos k_y a, \quad (5.55)$$

with the other diagonal elements just interchanging  $x \leftrightarrow y$  indices, giving rise to obvious square symmetry.

## VI. COMMENTS OF FITTING OF PARAMETERS

The tight binding hopping parameters (and overlap parameters, if used) are commonly fit to an independently calculated band structure. Denote the parameters by a vector  $\vec{t}$  and the set of eigenvalues to be fit by  $\{\varepsilon_K\}$ , where  $K$  will include both  $\vec{k}$  and band indices. The mathematical problem is to minimize the “residual”  $\mathcal{R}$

$$\delta\mathcal{R} = \delta \sum_K g_K [E_K(\vec{t}) - \varepsilon_K]^2 = 0. \quad (6.56)$$

Here  $g_K$  is a weight to be chosen as desired (for example, to fit eigenvalues around the Fermi energy more closely than elsewhere), and  $\{E_K\}$  are the eigenvalues resulting from the TB secular equation, which depend on the parameters.

Carrying out the minimization with respect to  $\vec{t}$  gives

$$\sum_K g_K (E_K(\vec{t}) - \varepsilon_K) \nabla_{\vec{t}} E_K(\vec{t}) = 0. \quad (6.57)$$

Linearizing around  $\vec{t} \approx \vec{t}_o$ :

$$\begin{aligned} \nabla_{\vec{t}} E_K(\vec{t}) &\approx \nabla_{\vec{t}} E_K(\vec{t}_o), \\ E_K(\vec{t}) &\approx E_K(\vec{t}_o) + (\vec{t} - \vec{t}_o) \cdot \nabla_{\vec{t}} E_K(\vec{t}_o) \end{aligned} \quad (6.58)$$



leads to

$$\sum_K g_k [E_K(\vec{t}_o) - \varepsilon_K + (\vec{t} - \vec{t}_o) \cdot \nabla_t (E_K(\vec{t}_o))] \nabla_t (E_K(\vec{t}_o)) = 0, \quad (6.59)$$

which can be written in a matrix form

$$\vec{b} + (\vec{t} - \vec{t}_o) \cdot A = 0. \quad (6.60)$$

This equation can be transposed to

$$\vec{t} = \vec{t}_o - \vec{b} \cdot A^{-1}, \quad (6.61)$$

where  $b$  and the matrix  $A$  are defined by

$$\begin{aligned} \vec{b} &= \sum_K g_k E_K(\vec{t}_o) - \varepsilon_K \nabla_t E_K(\vec{t}_o), \\ A_{p,q} &= \sum_K g_k [\nabla_t E_K(\vec{t}_o)]_p [\nabla_t E_K(\vec{t}_o)]_q. \end{aligned} \quad (6.62)$$

The Hellman-Feynman theorem gives a simple method of calculating the derivatives. Given the matrix eigenvalue equation

$$\hat{H}(\vec{k}) |k\alpha, \vec{t}\rangle = E_{\vec{k},\alpha} |k\alpha, \vec{t}\rangle \quad (6.63)$$

where  $\alpha$  is the band index, the derivative is

$$\nabla_t E_{\vec{k},\alpha}(\vec{t}) = \langle k\alpha, \vec{t} | (\nabla_t \hat{H}(\vec{k})) | k\alpha, \vec{t} \rangle. \quad (6.64)$$

Since the Hamiltonian matrix elements depend linearly on the components of  $\vec{t}$ , the derivative in this last equation is trivial and can be evaluated exactly by finite difference (numerically; of course, if the matrix is in analytic form it can be obtained trivially). A note to programmers: for Nb, where a  $9 \times 9$  matrix was used, with 31 parameters and 55  $\vec{k}$  points, there are

$$\frac{9 \times 10}{2} \times 55 \times 31 = 76,725 \quad (6.65)$$

such derivatives!

Eq. 6.61 provides an iterative procedure for fitting the parameters. Like all such schemes, it will converge if the starting point is sufficiently close to a good minimum. There will in general be many local minima, and the criterion for a best fit may be somewhat subjective as well. However, some study of these problems (unpublished) seems to indicate that non-uniqueness of the fit is not a big problem. Although a converged result is certainly not unique, restarting the iteration procedure in practice leads to sets of parameters whose differences have little physical significance, *i.e.* the large parameters are fairly well determined.

- 
- [1] W. E. Pickett and P. B. Allen, Solid State Commun. (1973).
  - [2] The square root of a matrix is ambiguous. After diagonalization, one can take  $\pm$  any of the diagonal

elements. Surely the physical choice is to take the positive square root of each diagonal element, though it is unclear how much this choice has been tested.